

Available online at <http://www.harmanpublications.com>

## World Journal of Science and Research

Harman Publications. All rights reserved



### Research Article

### Biochemistry

#### IN VITRO ANTI- INFLAMMATORY STUDIES ON SILVER NANOPARTICLES SYNTHESIZED FROM *Azima tetraacantha* LEAF

Manimegalai B<sup>1</sup> and Velavan S<sup>2</sup>.

<sup>1</sup>Research Scholar, P.G and Research Department, Marudupandiyar College, Thanjavur,

<sup>2</sup>Associate Professor, P.G and Research Department, Marudupandiyar College, Thanjavur,

\*Corresponding author: mayavelvan@gmail.com

#### ABSTRACT

#### Article Info:

This paper presents an empirical analysis on the use of aqueous extract of *Azima tetraacantha* for the production of silver nanoparticles (AgNPs) from aqueous silver nitrate. The *Azima tetraacantha* extract, AgNPs and Diclofenac sodium at different concentrations was incubated with egg and Bovine serum albumin in controlled experimental conditions and subjected to determination of absorbance to assess the anti-inflammatory property. Diclofenac sodium was used as the reference drug. The present findings exhibited a concentration dependent inhibition of protein denaturation by the *Azima tetraacantha* extract and AgNPs. The effect of AgNPs was found to be high when compared with the *Azima tetraacantha* extract and near to the diclofenac sodium. From the present study it can be concluded that AgNPs possessed marked *in vitro* anti-inflammatory effect against the denaturation of protein.

**Citation:** Manimegalai. B and Velavan S (2016) *In vitro* anti- inflammatory studies on silver nanoparticles synthesized from *Azima tetraacantha* leaf. *World Journal of Science and Research*. 1(3): 84-87.

Received on 18<sup>th</sup> Sep. 2016

Accepted on 15<sup>th</sup> Sep. 2016

Online September 2016

#### Keywords:

Silver nanoparticles, *Azima tetraacantha* extract, Diclofenac sodium, Anti-inflammatory effect

### INTRODUCTION

Inflammation is the complex response of the immune system to infection and injury that leads to removal of ending factors and restoration of tissue structure and physiological function (Ricciotti E, FitzGerald, 2011). The symptoms of inflammation are characterized by pain, heat, redness, swelling and loss of function. It can be classified into two major types either acute or chronic, based on the duration of the inflammatory reaction. Tough initiated as a

protective phenomenon, loss of regulation of this complex process can lead to the development of various inflammatory disorders.

Herbs and herbal extracts have been used to treat various ailments since ages. Their derivatives have attracted tremendous attention therapeutically and are promising as remedies to treat diseases of diversified origin. Herbs especially have fallen into limelight, anticipating their replacement with sophisticated

drugs. More than 50% of modern drugs existing in clinical use today are derived from plants. Metal nanoparticles have proved to be of significance due to their lesser volume to surface area ratio along with their catalytic, optical, electrical and magnetic characteristics (Nelson *et al.*, 2010), that are extensively used owing to their antimicrobial properties. Silver nanoparticles contribute even more, agents. Moreover, they are highly conductive, chemically stable and highly economical (Niraimathiet *et al.*, 2014). The plant extract was used for the preparation of silver nanoparticles owing to its least toxicity and lesser need for elaborate purification as compared to the chemical methods. The present work essentially deals with increasing therapeutic efficacy of the selected drug in its nanoparticle form. In the present study to synthesis of silver nanoparticles using the aqueous leaf extract of *Azimatetracantha* and evaluate the anti-inflammatory activity.

## MATERIALS AND METHODS

### Preparation of leaf extract

The dried leaves were pulverized well with mortar and pestle to make a powder. Twenty grams of powder sample was mixed into 100 ml of deionized water and the mixture was boiled for 10 min. After cooling the leaf extract was filtered with Whatman No. 1 filter paper. The filtrate was stored at 4°C for further use.

### Synthesis of Ag nanoparticles using leaf extracts

For the Ag nanoparticles synthesis, 5 ml of *Azimatetracantha* leaf extract was added to 45 ml of 1 mM aqueous AgNO<sub>3</sub> solution in a 250 ml Erlenmeyer flask. The flask was then incubated in the dark at 5hrs (to minimize the photo activation of silver nitrate), at room temperature. A control setup was also maintained without leaf extract. The Ag nanoparticle solution thus obtained was purified by repeated centrifugation at 10,000 rpm for 15 min followed by re-dispersion of the pellet in de-ionized water. Then the Ag nanoparticles were freeze dried using SEM analysis (Arunachalam *et al.*, 2012).

### IN VITRO ANTI-INFLAMMATORY ACTIVITY

*In vitro* anti-inflammatory activity was carried out by the method of Sangita Chandra *et al.* (2012). *In vitro* anti-inflammatory activity was carried out by the method of Sangita Chandra *et al.* (2012).

## RESULTS AND DISCUSSION:

There are certain problems in using animals in experimental pharmacological research, such as

ethical issues and the lack of rationale for their use when other suitable methods are available or could be investigated. Hence, in the present study the protein denaturation bioassay was selected for *in vitro* assessment of anti-inflammatory property of *Azimatetracantha* leaf extract and AgNPs. Denaturation of tissue proteins is one of the well-documented causes of inflammatory and arthritic diseases. Production of auto antigens in certain arthritic diseases may be due to denaturation of proteins (Egg albumin and Bovine serum albumin) *in vivo* (Opie, 1962; Umopathy *et al.*, 2010). Agents that can prevent protein denaturation therefore, would be worthwhile for anti-inflammatory drug development.


The use of nano-herbal-technology to synthesize compounds with improved anti-inflammatory properties is an area of current research by many scientists. In our study, we report the non-toxic, practical and environmentally benevolent approach for the synthesis of silver nanoparticles using the aqueous leaf extract of *Azimatetracantha* with potent anti-inflammatory activity. The synthesize and characterization of AgNPs from *Azimatetracantha* leaf extract showed the particle size between 10-80nm as well the cubic structure of the nanoparticles reported in our earlier report (Manimegalai and Velavan, 2015).

The increments in absorbances of test samples with respect to control indicated stabilization of protein i.e. inhibition of heat-induced protein denaturation by *Azimatetracantha* leaf extract, AgNPs and reference drug diclofenac sodium. The present findings exhibited a concentration dependent inhibition of protein denaturation by the *Azimatetracantha* leaf extract and AgNPs. The lowest activity of *Azimatetracantha* leaf extract, AgNPs and Diclofenac sodium were 10.32%, 14.56% and 21.03% in the concentration of 100µg/ml respectively while the highest activity of *Azimatetracantha* leaf extract, AgNPs and Diclofenac sodium were 68.67%, 82.64% and 89.65% in the concentration of 500µg/ml respectively. The greatest effect of AgNPs (500 µg/ml) was found to be near to standard diclofenac sodium. The half inhibition concentration (IC<sub>50</sub>) of *Azimatetracantha* leaf extract, AgNPs and diclofenac sodium were 359.22, 287.42µg/ml-1 and 237.14µg/ml-1 respectively. From the present study it can be concluded that AgNPs showed marked *in vitro* anti-inflammatory effect against the denaturation of protein (Table 1 and Figure 1). Our result agrees with the earlier report (Aparna Mani *et al.*, 2015; Giridharan *et al.*, 2014)

**Table 1: Effect of Azimatetracantha, AgNPs and Diclofenac sodium on protein denaturation (Fresh egg albumin)**

Groups	Concentrations	% of inhibition		
		<i>Azimatetracantha</i>	AgNPs	Diclofenac sodium (Standard)
Group I	100µg/ml	10.32±0.72	14.56±1.01	21.03±1.47
Group II	200µg/ml	35.74±2.50	42.65±2.98	51.65±3.61
Group III	300µg/ml	42.36±2.96	55.78±3.90	62.35±4.36
Group IV	400µg/ml	52.84±3.69	64.86±4.54	76.25±5.33
Group V	500µg/ml	68.97 ± 4.82	82.64±5.78	89.65±6.27
<b>IC<sub>50</sub> (µg/ml)</b>		359.22	287.42	237.14

Values are expressed as Mean ± SD for triplicates



**Figure 1: Effect of Azimatetracantha, AgNPs and Diclofenac sodium on protein denaturation (Fresh egg albumin)**


The present findings exhibited a concentration dependent inhibition of protein (Bovine serum albumin) denaturation by the *Azimatetracantha* leaf extract and AgNPs. The lowest activity of *Azimatetracantha* leaf extract, AgNPs and Diclofenac sodium were 20.56%, 25.36% and 23.75% in the concentration of 100µg/ml respectively while the highest activity of *Azimatetracantha* leaf extract, AgNPs and Diclofenac sodium were 71.85%, 86.24% and 91.52% in the concentration of 500µg/ml respectively. The half inhibition concentration (IC<sub>50</sub>)

of *Azimatetracantha* leaf extract, AgNPs and ascorbic acid were 281.75, 131.89µg/ml<sup>-1</sup> and 154.91µg/ml<sup>-1</sup> respectively. The greatest effect of AgNPs (500 µg/ml) was found to be near to standard diclofenac sodium. From the present study it can be concluded that AgNPs showed marked *in vitro* anti-inflammatory effect against the denaturation of protein (Table 2 and Figure 2). Our result agrees with the earlier report (Aparna Mani et al., 2015; Giridharan et al., 2014)

**Table 2: Effect of Azimatetracantha, AgNPs and Diclofenac sodium on protein denaturation (Bovine serum albumin)**

Groups	Concentrations	% of inhibition		
		<i>Azimatetracantha</i>	AgNPs	Diclofenac sodium (Standard)
Group I	100µg/ml	20.56±1.43	25.36±1.77	23.75±1.66
Group II	200µg/ml	43.65±3.05	49.65±3.47	58.65±4.10
Group III	300µg/ml	50.25±3.51	58.71±4.10	64.54±4.51
Group IV	400µg/ml	61.74±4.32	71.56±5.00	74.83±5.23
Group V	500µg/ml	71.85 ± 5.02	86.24±6.03	91.52±6.40
<b>IC<sub>50</sub> (µg/ml)</b>		281.75	131.89	154.91

Values are expressed as Mean ± SD for triplicates



**Figure 2: Effect of *Azimatetracantha*, AgNPs and Diclofenac sodium on protein denaturation (Bovine serum albumin)**

### CONCLUSION

The synthesised silvernanoparticles are capped by the phytochemicals of *Azimatetracantha* leaf extract especially flavonoids and show significant anti-inflammatory effects. In conclusion combining the benefits of phytomedicine with nanomedicine can result in the formation of more efficient silver nanoparticles. This finding suggests that the synthesis of AgNPs using *Azimatetracantha* leaf extract could be a good source for developing green nano-medicine for the management of inflammation.

### REFERENCES

Aparna Mani KM, Seethalakshmi S, Gopal V (2015) Evaluation of In-vitro Anti-Inflammatory Activity of Silver Nanoparticles Synthesised using *Piper Nigrum* Extract. *J Nanomed Nanotechnol* 6: 268; 1-5.

Arunachalama R, Sujatha Dhanasingha, Balasaraswathi Kalimuthua, Mani Uthirappana, Chellan Rosea., Asit Baran Mandal. (2012) Phytosynthesis of silver nanoparticles using *Coccinia grandis* leaf extract and its application in the photocatalytic degradation. *Colloids and Surfaces B: Biointerfaces* 94 226– 230.

Giridharan T., Chandran Masi, S. Sindhu and P. Arumugam. Studies on Green Synthesis, Characterization and Anti-proliferative Potential of Silver Nano Particle using *Dodonaea viscosa* and *Capparis decidua*. *Biosci., Biotech. Res. Asia*, 11(2), 665-673 (2014)

Manimegalai B And Velavan S. Green Synthesis Of Silver Nanoparticles Using *AzimaTetracantha* Leaf Extract And Evaluation Of Their Antibacterial And In

*Vitro* Antioxidant Activity. *Nanoscience And Nanotechnology: An International Journal* 2015; 5(2): 9-16.

Nelson Duran, Priscyl D. Marcato, Roseil De Conti, Oswaldo L. Alves, Fabio T. M. (2010) Costa and Marcelo Brocchi Potential Use of Silver Nanoparticles on Pathogenic Bacteria, their Toxicity and Possible Mechanisms of Action *J. Braz. Chem. Soc.*; (21)949.

Niraimathi K. L., R. Lavanya, V. Sudha, P. Brindha (2014) Green Synthesis And Characterization of Silver Nanoparticles From Aqueous Extract of *Basella alba* And Their In-Vitro Antioxidant Potentials. *International Journal of Pharmacy and Pharmaceutical Sciences*; (6) 10

Opie EL. (1962) On the relation of necrosis and inflammation to denaturation of proteins. *J Exp Med*; 115: 597-608.

Ricciotti E, FitzGerald GA (2011) Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol* 31: 986-1000.

Sangita Chandra, Priyanka Chatterjee, Protapaditya Dey and Sanjib Bhattacharya. (2012) Evaluation of *in vitro* anti-inflammatory activity of coffee against the denaturation of protein. *Asian Pacific Journal of Tropical Biomedicine*, 178-180.

Umamathy E, Ndebia EJ, Meeme A, Adam B, Menziwa P, Nkeh-Chungag BN, et al. An experimental evaluation of *Albica setosa* aqueous extract on membrane stabilization, protein denaturation and white blood cell migration during acute inflammation. *J Med Plants Res* 2010; 4: 789-795.