




Available online at <http://www.harmanpublications.com>

**World Journal of Science and Research**

Harman Publications. All rights reserved



**Research Article**

**Siddha Medicine**

**CHARACTERIZATION OF BIOACTIVE COMPOUNDS IN ALCOHOLIC EXTRACT OF *Hemidimus indicus* and *Alpenia officinarum* USING GC-MS TECHNIQUE**

**G. Vijayabaskar and V. Elango\***

Department of Siddha Medicine, Tamil University, Thanjavur, Tamil Nadu, India

**ABSTRACT**

Plants have been an important source of medicine with qualities for thousands of years. Phytochemicals are the chemicals extracted from plants. These organic chemicals are classified as primary or secondary constituents, depending on their role in plant metabolism. GC-MS method used for the analysis of the obtained extract can be an interesting tool for testing the amount of some active principles in herbs used in various industries. The aim of this study was to carry out for identification of bioactive compounds from the alcoholic extract of *Hemidimus indicus* and *Alpenia officinarum* by Gas chromatography and Mass spectroscopy (GC-MS). GCMS analysis of alcoholic extract was done by standard protocol using the equipment Perkin-Elmer Gas Chromatography–Mass Spectrometry, while the mass spectra of the compounds found in the extract was matched with the National Institute of Standards and Technology (NIST) library. The GC-MS analysis revealed the presence of various compounds like Seventeen (17) compounds were identified in *Hemidimus indicus* where and Fourteen (14) compounds were identified in *Alpenia officinarum*. In the alcoholic extract of *Hemidimus indicus* and *Alpenia officinarum* These findings support the traditional use of *Hemidimus indicus* and *Alpenia officinarum* in various disorders.

**Citation:** G. Vijayabaskar and V. Elango. (2018). Characterization of bioactive compounds in alcoholic extract of *Hemidimus indicus* and *Alpenia officinarum* using GC-MS technique, *World Journal of Science and Research*. 3(2): 16-22 (2018).

**Article Info:**

*Received on 12<sup>th</sup> May 2018*

*Accepted on 22<sup>th</sup> June 2018*

*Online July 2018*

**Keywords:**

Gas chromatography and Mass spectroscopy, *Hemidimus indicus* and *Alpenia officinarum* Phytochemistry.

**\*Corresponding author**

**V. Elango,**  
Department of Siddha Medicine, Tamil University, Thanjavur, Tamil Nadu, India

## INTRODUCTION

Plants have been an important source of medicine with qualities for thousands of years. Plants are used medicinally in different countries, and they are the source of many potent and powerful drugs. Mainly on traditional remedies such as herbs for their history, they have been used as popular folk medicines (Sathyaprabha *et al.*, 2010). It has been shown that *in vitro* screening methods could provide the needed preliminary observations necessary to elect crude plant extracts with potentially useful properties for further chemical and pharmacological investigations (Mathekaga and Meyer, 1998).

Phytochemistry or plant chemistry has developed in recent years as a distinct discipline, somewhere in between natural product organic chemistry and plant biochemistry and is closely related to both. It is concerned with the enormous variety of organic substances that are elaborated with and accumulated by plants and deals with the chemical structures of these substances, their biosynthesis, turn over and metabolism, their natural distribution and their biological function (Harborne, 1986).

Phytochemicals are the chemicals extracted from plants. These organic chemicals are classified as primary or secondary constituents, depending on their role in plant metabolism. Primary constituents include the common sugars, aminoacids, proteins, purines and pyrimidines of nucleic acids, chlorophyll's etc. Secondary constituents are the remaining plant chemicals such as alkaloids (derived from aminoacids), terpenes (a group of lipids) and phenolics (derived from carbohydrates) (Liu, 2004). Plant produces these chemicals to protect itself but recent research demonstrates that emphasizes the plant source of most of these protective, disease-preventing compounds. A true nutritional role for phytochemicals is becoming more probable every day as research uncovers more of their remarkable benefits (Hamburger M and Hostettmann, 1991). Within a decade, there were a number of dramatic advances in analytical techniques including TLC, UV, NMR and GC-MS that were powerful tools for separation, identification and structural determination of phytochemicals (Roberts, Xia, 1995).

Gas Chromatography Mass Spectroscopy (GC-MS) a hyphenated system which is a very compatible technique and the most commonly used technique for the identification and quantification of biochemical components of medicinal plants (Ronald Hites, 1997). The chosen medicinal plant namely as *Hemidimus indicus* and *Alpenia*

*officinarum* belongs. *Hemidimus indicus* and *Alpenia officinarum* is widely distributed in India, Nepal and Bhutan. The aim of this study is to determine the organic compounds present in the *Hemidimus indicus* and *Alpenia officinarum* extract with the aid of GC-MS Technique.

## MATERIALS AND METHODS

### Plant material and preparation of extracts:

The roots *Hemidimus indicus* and *Alpenia officinarum* were purchased from Traditional Medicinal shop, Thanjavur, Tamil Nadu, India. Healthy roots were washed several times with distilled water to remove the traces of impurities from the roots. Shade dried at room temperature for about 10 days and ground in to fine powder using mechanical grinder. The powder was extracted with alcohol. A semi solid extract was obtained after complete elimination of alcohol under reduced pressure. The *Hemidimus indicus* and *Alpenia officinarum* root extracts were stored in refrigerator until used.

### GC –MS analysis

GC-MS analysis was carried out on a GC clarus 500 Perkin Elmer system comprising a AOC-20i autosampler and gas chromatograph interfaced to a mass spectrometer instrument employing the following conditions: column Elite-1 fused silica capillary column (30 x 0.25mm ID x 1µMdf, composed of 100% Dimethyl polydioxane), operating in electron impact mode at 70eV; Helium gas (99.999%) was used as carrier gas at a constant flow of 1 ml /min and an injection volume of 0.5 µl was employed (split ratio of 10:1) injector temperature 250 °C; ion-source temperature 280 °C. The oven temperature was programmed from 110 °C (isothermal for 2 min), with an increase of 10 °C/min, to 200°C, then 5°C/min to 280°C, ending with a 9min isothermal at 280°C. Mass spectra were taken at 70eV; a scan interval of 0.5 seconds and fragments from 40 to 450 Da. Total GC running time is 36min. min. The relative percentage amount of each component was calculated by comparing its average peak area to the total areas. Software adopted to handle mass spectra and chromatograms was a TurboMass Ver 5.2.0.

## RESULTS AND DISCUSSION

Gas chromatography – mass spectrometry (GC-MS) is a method that combines the features of gas-liquid chromatography and mass spectrometry to identify different substances within a test sample (Kell *et al.*, 2005). In the last few years, GC-MS has become firmly established as a key technological platform for secondary metabolite profiling in both plant and non-plant species (Fernie

*et al.*, 2004). Plants have an almost limitless ability are phenols or their oxygen substituted derivatives. Most are secondary metabolites, of which at least 12,000 have been isolated, a number estimated to be less than 10% of the total. These substances serve as plant defense mechanisms against, insects and herbivores. Flavonoids exhibit several biological effects such as anti-inflammatory, anti-fungal, anti-hepatotoxic and anti-ulcer actions (De-Fatima *et al.*, 2006). Interpretation on mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology (NIST) having more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library. The name, molecular weight and structure of the components of the test materials were ascertained.

Seventeen (17) compounds were identified in *Hemidimus indicus* by GC-MS analysis. The active principles with their retention time (RT), molecular formula, molecular weight (MW) and concentration (%) are presented in (Table 1 and Fig 1). The prevailing compounds were Hexadecanoic acid, ethyl ester, Benzaldehyde, 2-hydroxy-4-methoxy and Linoleic acid ethyl ester. The biological activity of selected compounds represent in Table 2.

Fourteen (14) compounds were identified in *Alpenia officinarum* by GC-MS analysis. The active principles with their retention time (RT), molecular formula, molecular weight (MW) and concentration (%) are presented in Table 3 and Fig 2. The biological activity of selected compounds represent in Table 4. The biological activities of identified compounds were listed (Table 2) are based on Dr.Duke's Phytochemical and

to synthesize aromatic substances, most of which Ethnobotanical Databases by Dr. Jim Duke of the Agricultural Research Service/USDA.

Among the identified phytochemicals hexadecanoic acid is suggested to be a fatty acid ester and it may employed as antioxidant, antimicrobial, flavor, hypocholesterolemic agent and larvicidal activities (Bodoprost and Rosemeyer, 2007; Falodun *et al.*, 2009). 1, 2-benzenedicarboxylic acid, diisooctyl ester is a plasticizer compound and acts as antimicrobial and antifouling agent (Heinonen *et al.*, 1998).

Compounds like n-hexadecanoic acid, 12-octadecanoic acid, dodecanoic acid, tetradecanoic acid, 1,2-Benzenedicarboxylic acid, dibutyl ester, hexadecanoic acid, ethyl ester and 9,12-octadecadienoic acid (Z,Z) were identified in the ethanolic leaf extract of *Vitex altissima*, a Verbenaceae member (Sathish *et al.*, 2012). Likewise, hexadecane, dodecanoic acid, nonadecane, eicosane, tetradecanoic acid, oleic acid, heptacosane, 9,12- octadecenoic acid, ethyl ester; n-hexadecanoic acid; 1,2-benzenedicarboxylic acid and 9-octadecenoic acid (Z)-ethyl ester were reported in *Clerodendrum inerme* and *C. phlomidis* leaves (Anandhi and Ushadevi, 2013; Balaji and Kilimozhi, 2014).

The investigation concluded that the stronger extraction capacity of alcoholic could have been produced number of active constituents responsible for many biological activities. So that those might be utilized for the development of traditional medicines and further investigation needs to elute novel active compounds from the medicinal plants which may be created a new way to treat many incurable diseases.




Fig.1: Compounds identified in the *Hemidimus indicus*

**Table.1: Compounds identified in the *Hemidimus indicus***

S.No.	RT	Name of the compound	Molecular Formula	Molecular Weight	Peak Area %
1.	5.15	exo-2,7,7-trimethylbicyclo[2.2.1]heptan-2-ol	C <sub>10</sub> H <sub>18</sub> O	154	0.27
2.	6.02	Benzenepropanol, $\alpha$ -methyl- $\beta$ -nitro-, (R*,R*)-(.-.-)-	C <sub>10</sub> H <sub>13</sub> NO <sub>3</sub>	195	0.41
3.	7.35	Benzaldehyde, 2-hydroxy-4-methoxy-	C <sub>8</sub> H <sub>8</sub> O <sub>3</sub>	152	19.75
4.	9.89	10,12-Octadecadiynoic acid	C <sub>18</sub> H <sub>28</sub> O <sub>2</sub>	276	0.73
5.	14.55	7,9-Di-tert-butyl-1-oxaspiro(4,5)deca-6,9-diene-2,8-dione	C <sub>17</sub> H <sub>24</sub> O <sub>3</sub>	276	0.34
6.	15.61	Hexadecanoic acid, ethyl ester	C <sub>18</sub> H <sub>36</sub> O <sub>2</sub>	284	0.40
7.	17.66	Androstan-17-one, 3-ethyl-3-hydroxy-, (5 $\alpha$ )-	C <sub>21</sub> H <sub>34</sub> O <sub>2</sub>	318	5.14
8.	17.88	Linoleic acid ethyl ester	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	308	6.88
9.	20.11	Methyl 5,7-hexadecadiynoate	C <sub>17</sub> H <sub>26</sub> O <sub>2</sub>	262	4.81
10.	21.69	10,13-Octadecadiynoic acid, methyl ester	C <sub>19</sub> H <sub>30</sub> O <sub>2</sub>	290	1.34
11.	26.07	Doconexent	C <sub>22</sub> H <sub>32</sub> O <sub>2</sub>	328	0.36
12.	27.16	9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)-	C <sub>21</sub> H <sub>36</sub> O <sub>4</sub>	352	1.25
13.	35.89	Cedran-diol, (8S,14)-	C <sub>15</sub> H <sub>26</sub> O <sub>2</sub>	238	3.16
14.	37.28	Lupeol	C <sub>30</sub> H <sub>50</sub> O	426	2.03
15.	38.16	Methyl 3-cis,9-cis,12-cis-octadecatrienoate	C <sub>19</sub> H <sub>32</sub> O <sub>2</sub>	292	11.83
16.	39.49	$\beta$ -Amyrin	C <sub>30</sub> H <sub>50</sub> O	426	38.67
17.	39.83	Fenretinide	C <sub>26</sub> H <sub>33</sub> NO <sub>2</sub>	391	2.63

**Table.2: Biological activity of phytochemicals identified in the alcoholic extract of *Hemidimus indicus***

S.No	Compounds name	Biological Active compounds**
1.	Hexadecanoic acid, ethyl ester	Nematicide, Pesticide, Lubricant, Antiandrogenic, Flavor, Hemolytic 5-Alpha reductase inhibitor Antimicrobial
2.	Benzaldehyde, 2-hydroxy-4-methoxy-	Antimicrobial
3.	Linoleic acid ethyl ester	Antiinflammatory, Hypocholesterolemic, Cancer preventive Hepatoprotective Nematicide, Insectifuge Antihistaminic, Antieczemic Antiacne, 5-Alpha reductase inhibitor Antiandrogenic, Antiarthritic, Anticoronary, Insectifuge

\*\*Duke's. Phytochemical and Ethnobotanical Databases, www.ars-gov/cgi-bin/duke/, 2013.




Fig.2: Compounds identified in the *Alpenia officinarum*

Table.3: Compounds identified in the *Alpenia officinarum*

S.No.	RT	Name of the compound	Molecular Formula	Molecular Weight	Peak Area %
1.	4.70	Benzenepropanal	C <sub>9</sub> H <sub>10</sub> O	134	0.67
2.	5.15	α-Terpineol	C <sub>10</sub> H <sub>18</sub> O	154	0.53
3.	5.92	2-Butanone, 4-phenyl-	C <sub>10</sub> H <sub>12</sub> O	148	2.97
4.	9.89	β-copaene	C <sub>15</sub> H <sub>24</sub>	204	0.37
5.	10.69	4-((1E)-3-Hydroxy-1-propenyl)-2-methoxyphenol	C <sub>10</sub> H <sub>12</sub> O <sub>3</sub>	180	1.60
6.	18.27	2(1H)Naphthalenone, 3,5,6,7,8,8a hexahydro-4,8a-dimethyl-6-(1-methylethenyl)-	C <sub>15</sub> H <sub>22</sub> O	218	6.35
7.	19.22	Alloaromadendrene oxide-(2)	C <sub>15</sub> H <sub>24</sub> O	220	5.71
8.	20.10	1H-Imidazole, 4,5-dihydro-2-(phenylmethyl)-	C <sub>10</sub> H <sub>12</sub> N <sub>2</sub>	160	28.72
9.	20.67	3-Oxo-5-phenylpentanoic acid	C <sub>11</sub> H <sub>12</sub> O <sub>3</sub>	192	1.13
10.	21.62	α-Benzyl malanohydroxamic acid hydrate	C <sub>10</sub> H <sub>12</sub> N <sub>2</sub> O <sub>4</sub>	224	16.50
11.	24.81	3-Isopropylidene tricyclo[4.3.1.1(2,5)]undecan-10-one	C <sub>14</sub> H <sub>20</sub> O	204	2.60
12.	26.01	Phenol, 2-methoxy-4-propyl-	C <sub>10</sub> H <sub>14</sub> O <sub>2</sub>	166	28.87
13.	26.54	trans-Dehydroandrosterone, trifluoroacetate	C <sub>21</sub> H <sub>27</sub> F <sub>3</sub> O <sub>3</sub>	384	2.79
14.	27.64	2,5-Octadecadiynoic acid, methyl ester	C <sub>19</sub> H <sub>30</sub> O <sub>2</sub>	290	1.18

**Table.3: Biological activities of bioactive compounds in *Alpenia officinarum***

S.No.	RT	Name of the compound	Biological Active compounds**
1	5.15	$\alpha$ -Terpineol	Anticonvulsant Activity, Antitumour activity, Anti-inflammatory activity Antioxidant and Antimicrobial Activity
2	9.89	$\beta$ -copaene	Antimicrobial Activity; Antioxidant Activity
3	26.01	Phenol, 2-methoxy-4-propyl-	Antimicrobial activity
4	26.54	trans-Dehydroandrosterone, trifluoroacetate	Steroid hormone like activity

\*\*Duke's. Phytochemical and Ethnobotanical Databases, www.ars-gov/cgi-bin/duke/, 2013.

## CONCLUSION

The investigation concluded that the stronger extraction capacity of alcoholic could have been produced number of active constituents responsible for many biological activities. So that those might be utilized for the development of traditional medicines and further investigation needs to elute novel active compounds from the medicinal plants which may be created a new way to treat many incurable diseases.

## REFERENCES

- Anandhi K and Ushadevi T. (2013) Analysis of phytochemical constituents and antibacterial activities of *Clerodendrum inerme* L. against some selected pathogens. *International Journal of Biotechnology and Allied Fields*, 1(7): 387-393.
- Balaji K and Kilimozhi D. (2014) GC-MS analysis of various extracts of *Clerodendrum phlomidis* leaf. Responsible responsible for many biological activities and its beneficial effects could be utilized to create an *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(1): 226-232.
- Bodoprost J and Rosemeyer H. (2007) Analysis of phenacyl ester derivatives of fatty acids from human skin surface by reversed-phase HPTLC: Chromatography mobility as a function of physicochemical properties. *International Journal of Molecular Sciences*, 8: 1111-1124.
- De-Fatima A, Modolo LV, Conegero LS, Pilli RA, Ferreira CV, Kohn LK and de-Carvalho JE. (2006) Lactones and their derivatives: biological activities, mechanisms of action and potential leads for drug design. *Curr. Med. Chem.* 13: 3371-3384.
- Falodun A, Siraj R and Choudary MI. (2009) GC-MS analysis of insecticidal leaf essential oil of *Pyrenacanthastaudtii* Hutch and Dalz (Icacinaceae). *Tropical Journal of Pharmaceutical Research*, 8: 139-143.
- Fernie AR, Trethewey RN, Krotzky AJ and Willmitzer L. (2004) Metabolite profiling: From diagnostics to systems biology. *Nat Rev Mol Cell Biol.*; 5:763-9.
- Hamburger M and Hostettmann, K. (1991) Bioactivity in plants: the link between phytochemistry and medicine. *Phytochemistry*. 30; 3864-74.
- Harborne, JB. (1986) Plant flavonoids in biology and medicine: Biochemical pharmacological, and structure-activity relationships. NY, USA: Alan R. Liss. pp. 15-24.
- Heinonen OP, Alnanes D and Virtamo T. (1998) Prostate cancer and supplementation with alpha-tocopherol and beta-carotene: incidence and mortality in a controlled trial. *Journal of the National Cancer Institute*, 90(6): 440-446.
- Kell DB, Brown M, Davey HM, Dunn WB, Spasic I and Oliver SG. (2005) Metabolic footprinting and systems biology: The medium is the message. *Nat Rev Microbiol*; 3:557-65

- Liu RH. (2004) Potential synergy of phytochemicals in cancer prevention: Mechanism of action. *Journal of Nutrition*, 134(12 Suppl.); 3479S–3485S.
- Mathekaga, AD and Meyer JJM. (1998) Antibacterial activity of South African *Helichrysum* species. *South Afr J Bot*; 64:293-5.
- Roberts JKM and Xia JH. (1995) High-resolution NMR methods for study of higher plants, *Methods Cell Biol.* 49; 245–258.
- Ronald Hites A. (1997) Gas Chromatography Mass Spectroscopy: Handbook of Instrumental Techniques for Analytical Chemistry, 609-611.
- Sathish SS, Janakiraman N and Johnson M. (2012) Phytochemical analysis of *Vitexaltissima*L. Using UV-VIS, FTIR and GC- MS. *International Journal of Pharmaceutical Sciences and Drug Research*, 4(1): 56-62.
- Sathyaprabha G, Kumaravel S, Ruffina D and Praveenkumar PA. (2010) comparative study on antioxidant, proximate analysis, antimicrobial activity and phytochemical analysis of *Aloe vera* and *Cissus quadrangularis* by GC-MS. *J Pharma Res*;3:2970-3.